

After retinal detachment surgery (no posturing required)

This leaflet is for patients who have undergone retinal detachment surgery and gives advice on what to do once you get home to aid healing.

Instructions to follow after your operation


When you go home after surgery, you should follow the following advice for approximately 4-6 weeks.

- You can carry out most daily activities as normal but it may take a few days to fully recover from the anaesthetic so it is a good idea to take things easy.
- 2. You may read or watch television, as you feel able.
- 3. It takes time for your vision to settle after your operation approximately 2-3 months. Please don't worry if you cannot see well immediately after the operation.
- 4. Use your eye drops and any other medication/tablets prescribed by the doctor.
- 5. Avoid rubbing or pressing on your eye.
- 6. Your eye may look red and be slightly painful after the operation. This will gradually get better as you use your eye drops and take any pain relief you have been prescribed. If this does not control the pain, contact the hospital on the number listed on the back of this leaflet.
- 7. After you have been for your first follow-up appointment and the doctor is happy with your progress, you can return to your normal routine. This generally means that you can return to work. You may take exercise. The only precaution you must take is to avoid direct injury to the eye. If you have any particular queries, e.g. resuming driving, please ask the doctor or nurse at the clinic.
- 8. If you notice your sight getting worse rather than better, you should contact the Eye Unit as soon as possible.

| Compassionate | Aspirational | Resourceful | Excellent |
|---------------|--------------|-------------|-----------|
| Compassionate | Aspirational | Resourceful | EXCELLETT |

Putting in eye drops

- 1. Wash your hands thoroughly.
- 2. Tilt your head back and look at the ceiling.
- 3. Gently pull the lower eyelid down until there is a small pocket (use a tissue or cosmetic pad to do this).



- 4. Squeeze the bottom of the dropper/bottle to release a drop into the eye.
- 5. Release the lower lid and close your eye for 30 seconds. Dab any excess fluid with a tissue.
- 6. Wash your hands.

Contacting us

If you have a minor eye problem, please seek advice from your GP, optician or pharmacist. If urgent, please attend Eye Casualty or call 111.

| Eye Casualty (Reading): | Mon-Fri 8.30am to 5pm; Sat & Sun & bank holidays 9am-12.30pm; Closed Christmas Day and New Year's Day. | |
|-----------------------------|--|--|
| Eye Casualty: Prince | Mon-Fri 8.30am to 5pm; Sat 8.30am- | |
| Charles Eye Unit (Windsor): | 12.30pm; Closed Sun & bank holidays. | |
| Dorrell Ward (Reading): | 0118 322 7172 | |
| Dorrell Ward (Reading). | (24 hours a day) | |
| Eve Day Unit (Peeding): | 0118 322 7123 | |
| Eye Day Unit (Reading): | (Mon-Fri 7am to 6pm) | |

To find out more about our Trust visit www.royalberkshire.nhs.uk

Please ask if you need this information in another language or format.

RBFT Ophthalmology, October 2023

Next review due: October 2025

| Compassionate | Aspirational | Resourceful | Excellent |
|---------------|--------------|-------------|-----------|
| | | | |