



Wisdom teeth removal

This leaflet is designed to give you more information about what the issues are with wisdom teeth, what treatment options are, where you may be referred to and what to expect if you are having treatment. Your dentist may also be able to discuss this with you.



Issues with lower wisdom teeth


Wisdom teeth are the last teeth to develop in your jaws. Often there is not enough space for them to come through into the correct position like the other teeth in your mouth so they can vary in their position. Please see the image (*right*) for their different positions. Some may not come through fully because they are 'impacted' against another tooth or may be partly covered by the gums (oral mucosa).


Some wisdom teeth may not appear to be causing a problem, but these (varied position, impaction and partially come through) and other issues may develop, which you may be unaware of.

There is strong evidence to show that wisdom teeth that are mesially (horizontally) positioned and have not erupted fully have a high risk of decay occurring not only in the wisdom tooth but also to the tooth in front of it.


If left alone the decay may get to the extent that you may develop pain from both teeth. If the tooth in front is decayed it will need to be repaired by the dentist after the wisdom tooth has been extracted.

The decay may be so bad that the tooth in front has to be removed as well. The decay may also spread to the bone around these teeth causing periodontal disease. Sometimes cysts can form around the wisdom tooth. A cyst occurs when fluid fills the sac that normally surrounds a developing wisdom tooth and can slowly increase in size.





A horizontal wisdom impacted teeth. There is early decay in the tooth in front. Your dentist may be able to repair this tooth.



A horizontal impacted wisdom tooth. There is extensive decay in the tooth in front. Both teeth will need to be removed.

If you have a medical issue that requires you to take medicines such as immune-suppressants or medicines that affect bone it is advisable to have an assessment of your teeth, including your wisdom teeth, before you start these medicines. If you were to have a tooth extracted while on these medicines this may slow down healing of a tooth socket. In rare cases the tooth socket may not heal at all. The assessment may recommend that the teeth which are badly decayed, including your wisdom teeth, should be removed.

If you are intending to travel abroad for a long length of time and access to a dentist or a specialist is difficult, you should consider having treatment for your wisdom teeth and other teeth before you leave.

Do I need to be referred?

Some wisdom teeth can be straightforward to remove. Your dentist might be able to remove them or may refer you to see a clinician who removes them routinely. You may need to go to another dental practice or a clinic in the community.

For wisdom teeth that are difficult to remove or there are medical issues or if you wish to see a specialist, you may need to be referred to the hospital. The specialist will often be a consultant in oral and maxillofacial surgery or an oral surgeon. They will have the experience and qualifications to provide you with the information you need to decide what you would like to do about your wisdom teeth. Often the first appointment will be a consultation only and if necessary another appointment will be needed to have your treatment.

What treatment options may be discussed?

- **No treatment but kept under review:** If the wisdom tooth is in a position where it has not been causing problems and there is a low risk of a problem developing, then an option can be for the wisdom tooth to be left where it is. Your dentist can continue to review your wisdom teeth along with the rest of your teeth when you attend for your routine check-up. When necessary your dentist may need to take an X-ray of the wisdom teeth. However, over time circumstances may change and the risk of a problem with your wisdom teeth may increase, and treatment will then be needed. The time of when treatment is needed is a decision that can be made with you and the dentist or the specialist.

It is important to keep your wisdom teeth under review as problems may develop without you knowing about them until severe symptoms develop.

- **Extraction (removal) treatment:** In most cases where a problem has developed or there is a high risk of a problem developing, the wisdom tooth could be removed completely or part removed. During your consultation these options will be discussed with you in detail along with what is involved and the risks of the procedure.


What does the complete removal of lower wisdom teeth involve?

If the wisdom tooth has not fully erupted into the mouth it is often necessary to make a cut in the gum over the tooth. Sometimes it is also necessary to remove some bone surrounding the wisdom tooth. This is all done under local anaesthetic, so you won't feel any pain during the procedure. The tooth may need to be cut into 2 or 3 pieces to remove it. Once the wisdom tooth has been removed the gum is put back into place with stitches. In the majority of cases these stitches are dissolvable and take around two weeks to disappear.

What are the main risks of removing a wisdom tooth?

Most patients will recover fully with no problems.

- **Bleeding:** There may be a little bleeding at the time of the extraction. This usually stops very quickly and is unlikely to be a problem. If you are at home and the area bleeds again this can usually be stopped by applying pressure over the area for at least 10 minutes with a rolled up handkerchief or swab. If the bleeding does not stop, please contact the department where you had your operation.
- **Altered sensation to the chin, lower lip or tongue:** Two nerves lie quite close to the lower wisdom teeth. The nerves can vary in their proximity to wisdom teeth. See the image below. The lingual nerve provides sensation in your tongue while the inferior dental nerve gives sensation to the lower lip and skin overlying the chin. Occasionally, these nerves may be bruised during tooth removal. If this happens then you may have either numbness or pins and needles in the tongue, lower lip, skin over the chin, or any combination of these. If the wisdom tooth is away from the nerve the risk to the inferior dental nerve is less than 1% (1 in every 100). However, if the nerve is in very close contact the risk may be as high as 9% (9 in every 100). The risk to the lingual nerve is less than 0.5% (less than 1 in every 200).



Compassionate	Aspirational	Resourceful	Excellent
---------------	--------------	-------------	-----------

- It may take up to 18 months for this to recover. Rarely the nerve may not recover fully and you may be left with small patches of numbness to your lip or chin. Taste is rarely affected.
- **Infection:** An infection in the in the socket can occur or the blood clot from extraction socket has dissolved away which may result in a dry socket (a painful dental condition). The incidence of this happening is low – around 2% (2 out of every 100). It can be minimised by keeping the area clean by rinsing with water or a mouthwash, starting the day after your procedure. It is advisable to give up smoking, or at least avoid smoking for a few days after having the wisdom tooth removed. Smoking can increase the risk of an infection or a dry socket. It may also be necessary for you to have a course of antibiotics after the extraction
- **Weakening of the jaw:** In exceptional cases wisdom tooth removal can cause significant weakening of the jaw with the resultant possibility of jaw fracture. The incidence of this happening is extremely low and is less than 1% (less than 1 in 100).

Coronectomy treatment

A coronectomy may be a treatment option. It may be offered if the inferior dental nerve is in close contact to the wisdom tooth. In this technique only the upper portion of the wisdom tooth is removed leaving some of the tooth roots behind. This is to minimise the risk of damage to the nerve. However, there is less than a 3% (less than 3 in every 100) risk that you may need to have the remaining roots taken out at a later date or there may be delayed (slow) healing. At the consultation an assessment will be made to see if this option is appropriate for you.

How long does it take to remove a wisdom tooth?

This varies. Some wisdom teeth may take only a few minutes to remove. More difficult wisdom teeth that need to be cut into pieces to remove can take around 20 minutes to extract.

Is there much pain or swelling after the removal of wisdom teeth?

It is likely that there will be some discomfort and swelling both on the inside and outside of your mouth after surgery. This is usually worse for the first three days but it may take up to two weeks before all the soreness goes. You may also find that your jaw is stiff and may not be able to open fully. You will need to eat a soft diet, such as soups and mashed/pureed foods, for a week or so. There may be some bruising of the skin of your face that can take up to a fortnight to fade away. The application of an ice pack during the first 24 hours after the extraction may help. Do not apply the ice pack directly to your skin – wrap it in a towel first. The older you are having a wisdom tooth removed the longer it will take to recover.

What type of anaesthetic is used for surgery?

A number of options are available and depend on how difficult the wisdom tooth is to remove:

- **Local anaesthetic:** this is an injection into the gum surrounding the wisdom tooth, rather similar to that you may have had at your dentist for a filling. The injection takes a couple of minutes to numb the area and means that you will feel no pain while the wisdom tooth is removed.
- **Local anaesthetic and intravenous sedation:** in addition to a local anaesthetic injection you can be given an injection into your arm. This makes you feel relaxed and less aware of the

procedure. You are awake but patients frequently have little or no memory of the operation.

- **General anaesthetic:** it is usually possible to remove wisdom teeth under a “day case” general anaesthetic, i.e. although you are put to sleep completely you will be able to go home on the same day as surgery.

Is there anything else I need to do after the extractions?

It is important to keep the extraction sites as clean as possible for the first few weeks after surgery. It may be difficult to clean your teeth around the sites of the extraction because it is sore. If this is the case it is best to keep the area free from food debris by gently rinsing with a mouthwash or warm salt water after eating (dissolve a flat teaspoon of kitchen salt in a cup of warm water), starting on the day after surgery. We will give you a separate leaflet on looking after your extraction site.

Do I need to take any time off work?

Usually it will be necessary to take a few days off work and avoid strenuous exercise for this time. Depending on the type of anaesthetic used, you may well not be able to drive (24 hours after intravenous sedation and for 48 hours after a general anaesthetic).

Further information

Royal College of Surgeons of England. <http://www.rcseng.ac.uk/>

British Association of Oral and Maxillofacial Surgeons. <http://www.baoms.org.uk/>

NHS Stop Smoking Service. <http://smokefree.nhs.uk/>

British Association of Oral Surgeons. <http://www.baos.org.uk/>

Contacting us

Oral & Maxillofacial Department, Tel: 0118 322 7139 or email: rbb-tr.cat1@nhs.net

To find out more about our Trust visit www.royalberkshire.nhs.uk

Please ask if you need this information in another language or format.

RBFT Department of Oral & Maxillofacial Surgery, June 2024

Next review due: June 2026