



Macular holes

We hope this information will answer some of your questions about macular holes and surgery to repair them. If there is anything you do not understand, or if you have any concerns, please tell us, as it is important that you understand everything before you go ahead with surgery and sign a consent form.

What is a macular hole?

The back of the eye has a light-sensitive lining called the retina, similar to the film in a camera. Light is focused through the eye onto the retina, allowing us to see. The centre part of the retina is called the macula – it is here that light must be focused for us to see fine detail, to be able to read and to see in colour.

A macular hole is a small, circular gap, which opens up at the centre of the retina. This causes blurred vision and often-distorted vision, where straight lines or letters look wavy or bowed. There may also be a patch of missing vision at the centre.



Is this the same as age-related macular degeneration?

No, macular holes and macular degeneration are different conditions although they affect the same area of the eye. They can sometimes both be present in the same eye.

Why do macular holes happen?

We do not know why macular holes develop. They most often occur in people aged 60-80, and is twice as common in women as men. We are increasingly aware that it mainly affects slightly long-sighted people. Other causes of macular holes include severe trauma to the eye, being very short-sighted (myopic), those who have had a retinal detachment or because of longstanding swelling of the central retina (cystoid macular oedema).

What happens if it is left untreated?

If untreated, there is a small chance that some macular holes can close spontaneously, with improvement in vision. In the majority of patients, the central vision will gradually get worse to a level where the patient is unable to read even the largest print on an eye test chart.

The condition does not affect the peripheral (at the edges) vision, and so patients will not go completely blind from this condition.

Can I develop macular holes in my other eye?

Careful examination can assess the risk of developing a macular hole in the other eye. Your surgeon will tell you your risk but this may be from extremely unlikely to a 1 in 10 chance. It is very important to monitor for any changes in vision of the fellow eye, and report these to your eye specialist/family doctor/optician urgently.

What is the treatment and how successful is it?

A macular hole can often be repaired by an operation called a **vitrectomy, peel and gas**. If the hole has been present for less than a year, the operation will be successful in closing the hole in about 90% of cases. Of these, more than 70% will be able to see two or three lines more down a standard vision chart, compared to before the operation. Even if this degree of improvement does not occur, the vision is at least stabilised and many patients find that they have less distortion. In a minority of patients, the hole does not close despite surgery and the central vision can continue to deteriorate; however, a second operation

can still be successful in closing the hole. It is important to understand that return to completely normal vision is not possible.

If I opted for surgery, does it matter how long I have had the macular hole?

There is evidence that relatively early treatment (within months) gives a better outcome in terms of improvement in vision. Studies have shown, however, that vision improvement may be possible in some patients with long-standing macular holes.

What does the operation involve?

Macular hole surgery is a form of keyhole surgery performed under a microscope, using 3 small incisions (1-2 mm in size) in the white of the eye for insertion of very fine instruments. Firstly, the vitreous jelly is removed (vitrectomy), and then a very delicate layer (the inner limiting membrane) is carefully peeled off the surface of the retina around the hole to release the traction forces that keep the hole open.

The eye is then filled with a temporary gas bubble, which presses against the hole to help it seal. The bubble of gas blocks the vision whilst it is present, but slowly disappears over a period of about 2-8 weeks. We usually put small, dissolving stitches in the eye. At the end of the operation, we may put a pad or bandage over your eye to protect it and reduce swelling.

How long does the operation take and do I need to have a general anaesthetic?

Macular hole surgery usually takes 45-90 minutes and can be done with the patient awake (local anaesthetic), or asleep (general anaesthetic), often as a day case procedure. Most patients opt for a local anaesthetic, which involves a numbing injection around the eye so that you feel no pain during the operation; this is sometimes supplemented with medication to reduce anxiety (sedation).

If you would prefer a general anaesthetic, we will ask you not to eat or drink for several hours before we take you to the operating theatre.

After the operation

If you have discomfort, we suggest that you take a pain reliever, such as paracetamol or ibuprofen, every 4-6 hours. It is normal to feel itching, sticky eyelids and mild discomfort for a while after macular hole surgery. It is common for your eye to water. Occasionally, the area surrounding the eye can become bruised. Any discomfort should ease after 2-3 days. In most cases, your eye will take about 6 weeks to heal. Try to rest while your eye is healing.

We will give you eye drops to reduce any inflammation and to prevent infection. We will explain how and when you should use them. Please do not rub your eye.

Certain symptoms could mean that you need prompt treatment. Please contact the hospital immediately if you have any of the following:

- A lot of pain
- Loss of vision
- Increasing redness of the eye

Do I have to posture face down after the operation?

The aim of face down posturing is to keep the gas bubble in contact with the hole as much as possible to encourage it to close. Whether you are required to posture, and for how long, will depend on the size of the macular hole, and also the preferences of your surgeon. There is evidence that posturing improves the success rate for larger holes, but it may not be needed for smaller holes.

If you are asked to do some face down posturing, your head should be positioned so that the tip of your nose points straight down to the ground. This could be done sitting at a table, or lying flat on your stomach on a bed or sofa. You should try to remain in this position for up to 50 minutes in each hour, depending on your consultant's advice, for the duration advised (usually 2-5 days after the operation). A short break of 10-15 minutes can be taken every hour to allow eating, trips to the bathroom etc. Your surgical team will advise on aids that can make face down posturing easier to manage e.g. a horseshoe-shaped pillow or frame.

Please remember that if you are not able to posture then there is still a good chance that the hole will close successfully.

If face down posturing is not prescribed, you may be simply advised not to lie on your back for a period of two weeks after the surgery, with some surgeons recommending sleeping in a chair or at 45° in bed, supported by pillows for the same period.

Can I travel after macular hole surgery?

You must not fly or travel to high altitude on land whilst the gas bubble is still in the eye (2-8 weeks). If ignored, the bubble will expand at altitude, causing very high pressure resulting in severe pain and permanent loss of vision. In addition, if you need a general anaesthetic whilst gas is in your eye, then it is vital that you tell the anaesthetist this fact so they can avoid certain anaesthetic agents, which can cause similar expansion of the bubble. None of these exclusions apply once the gas has fully absorbed. You will notice the bubble shrinking and will be aware when it has completely gone but please check with your consultant if in doubt.

How much time will I need off work?

Most people will need two weeks off work. Your vision is reduced while the gas bubble is in the eye and this also affects depth perception. However, it depends on the type of work you do and the speed of recovery. You should discuss this with your surgeon.

What are the potential complications of macular hole surgery?

As with any procedure there may be risks involved and you should discuss these fully with the consultant involved prior to your operation, however it is unlikely that you will suffer harmful effects from a macular hole operation. In a small minority, the vision may end up worse than before the surgery, and there is even a tiny chance of total loss of sight. Six specific complications of macular hole surgery, which you must be aware of, are outlined below:

1. **Failure of the macular hole to close:** this occurs in 1-2 out of 10 patients. In most circumstances, it is possible to repeat the surgery. If the hole fails to close, then the vision may be a little worse than prior to the surgery.
2. **Cataract:** this means that the natural lens in eye has gone cloudy. If you have not already had a cataract operation, you will almost certainly get a cataract after the surgery, usually within a year but it can happen very rapidly. As cataract is inevitable, you may be offered combined surgery with cataract extraction at the same time as the macular hole repair.
3. **Retinal detachment:** the retina detaches from the back of the eye in 1-2% of patients undergoing macular hole surgery. The vast majority of retinal detachments are repairable, but further surgery is required and this can be a potentially blinding complication.
4. **Bleeding:** this occurs very rarely, but severe bleeding within the eye can result in blindness.
5. **Infection:** this is also very rare and would be expected to occur in about 1 in 1000 patients, but if it occurs needs further treatments and can lead to blindness.
6. **Raised eye pressure:** an increase in pressure within the eye is quite common in the days after macular hole surgery, usually due to the expanding gas bubble. In most cases, it is short-lived and controlled with extra eye drops and/or tablets to reduce the pressure, preventing any harm coming to the eye. If the high pressure is extreme or becomes prolonged, there may be some damage to the optic nerve as a consequence. In the majority, this damage does not adversely affect the vision, but some patients require long-term treatment to keep the eye pressure controlled.

Will I need to use eye drops or medication after surgery?

Three types of drops are usually prescribed after surgery: an antibiotic, a steroid and a pupil-dilating agent. Patients are seen again in the clinic about two weeks after the surgery. If all is well, then the drops are reduced over the following 2-4 weeks. If the eye pressure is raised

following surgery, additional drops and/or tablets may be prescribed to treat this.

When will I need to be seen again after the surgery?

Post-operative review is usually performed the next day (or within a few days) after surgery; then 10-14 days later and, provided all is well, about 3 months later.

Will I have to get my glasses changed?

Most people will need to change their spectacle prescription at some point after surgery. This would normally be at about 3-months following the operation, after the gas bubble has gone. As each case is different, please check with your surgeon before visiting an optician.

Where can I find more information?

- The RNIB have further information on macular holes, especially some practical advice: Helpline 0303 123 9999; website: www.rnib.org.uk; email helpline@rnib.org.uk
- The Macular Disease Society: Helpline 0845 241 2041; website: www.maculardisease.org; email: info@maculardisease.org

References

This information is based on a variety of sources, including latest research published in peer-reviewed scientific journals. It has also been scrutinized by a panel of experts from the Britain & Eire Association of Vitreoretinal Surgeons (BEAVRS). If you require further information about this, please ask your surgeon.

Contacting us

If you have a minor eye problem, please seek advice from your GP, optician or pharmacist. If urgent, please attend Eye Casualty or call 111.

| | |
|--|---|
| Eye Casualty (Reading): | Mon-Fri 8.30am to 5pm; Sat & Sun & bank holidays 9am-12.30pm; Closed Christmas Day and New Year's Day. |
| Eye Casualty: Prince Charles Eye Unit (Windsor): | Mon-Fri 8.30am to 5pm; Sat 8.30am-12.30pm; Closed Sun & bank holidays. |
| Dorrell Ward (Reading): | 0118 322 7172 (24 hours a day) |
| Eye Day Unit (Reading): | 0118 322 7123 (Mon-Fri 7am to 6pm) |
| Eye Day Unit (PCEU Windsor) | 01753 636496 Mon-Fri 7am to 6pm) |

Outside of Eye Casualty hours, you should telephone your GP's out of hours' service, ring NHS 111 or if you have serious concerns, visit your nearest Emergency Department (A&E).

To find out more about our Trust visit www.royalberkshire.nhs.uk

Please ask if you need this information in another language or format.

Ophthalmology, June 2023. Next review due: June 2025