

## Gardasil 9 suspension for injection in a pre-filled syringe Human Papillomavirus 9-valent Vaccine (Recombinant, adsorbed)

NHS number	
Name (in BLOCK capitals)	
Date of Birth (dd/mm/yyyy)	
Hospital MRN number	

**Please answer the following questions**

Have you had a Gardasil 9 vaccine previously?	Yes <input type="checkbox"/> No <input type="checkbox"/>
Do you have a fever or infection or feel unwell?	Yes <input type="checkbox"/> No <input type="checkbox"/>
Have you any known allergies including YEAST?	Yes <input type="checkbox"/> No <input type="checkbox"/>
Have you had an allergic reaction to any medication?	Yes <input type="checkbox"/> No <input type="checkbox"/>
Do you have any illness which may affect your immune system?	Yes <input type="checkbox"/> No <input type="checkbox"/>
Are you taking any medicines (e.g. steroids) which may affect your immune system?	Yes <input type="checkbox"/> No <input type="checkbox"/>
Do you have any blood disorders or take anti-clotting medication? e.g. Warfarin	Yes <input type="checkbox"/> No <input type="checkbox"/>
Have you experienced any significant health problems after previous vaccine administration?	Yes <input type="checkbox"/> No <input type="checkbox"/>
<b>If you answered 'Yes' to any of the above questions, please provide further detail below:</b>	

**What Gardasil 9 contains:**

The active substances are: highly purified non-infectious protein for each of the Human Papillomavirus types (6,11,16,18,31,33,45,52 and 58)  
 Human Papillomavirus = HPV  
 L1 protein in the form of virus like particles produced in yeast cells (*Saccharomyces cerevisiae* CANADE 3C-5 (Strain 1895) by recombinant DNA technology.  
 Adsorbed on amorphous aluminium hydroxyphosphate sulfate adjuvant (0.5 milligrams A1).  
 Amorphous aluminium hydroxyphosphate sulfate is included in the vaccine as an adjuvant. Adjuvants are included to improve the immune response.  
 The other ingredients in the vaccine suspension are: sodium chloride, histidine, polysorbate 80, borax and water for injections.

**If you are immunocompromised you may have a lower immune response to the vaccine.**

**In case of allergic reaction, patients are advised to be observed for 15 minutes after they receive the Gardasil 9 vaccine.**

**Gardasil 9 may slightly and temporarily affect the ability to drive or use machines.**

**THE GARDASIL 9 VACCINE CANNOT CAUSE HPV RELATED DISEASES.**

**Side effects**, if experienced are likely to occur within the first 24-48 hours post vaccination.

If you experience, pain, swelling and/or redness at the site of the injection a cold compress may help.

If you experience mild fever, headache, malaise, myalgia, arthralgia, analgesics such as Paracetamol may help.

**Immune response is generally induced within 2-3 weeks post vaccination. A booster dose is not yet determined.**

**Sign below to confirm you have read and understood the above information and consent to having the Gardasil 9 vaccination course.**

<b>Individual's signature:</b>		<b>Date:</b>	
<b>Vaccinator's name (CAPITALS):</b>		<b>Dose:</b>	0.5ml
		<b>Batch No:</b>	
		<b>Expiry date:</b>	
<b>Vaccinator's signature:</b>		<b>Injection site:</b>	Left / Right Deltoid