



This leaflet outlines 'Sick Day Rules' for children with diabetes who are on insulin pumps and their family.

When you are ill your body needs more energy to fight the infection.

Glucose is released from your body stores to do this, so the blood glucose tends to go up even if you are not eating.

This means that you need more insulin than usual when you are ill. If you do not give enough insulin then your body cannot use the extra glucose so will start to break down fat for energy.

When fat is broken down, this produces ketones.

Ketones are very dangerous!



What are ketones?

Ketones are acids which can quickly make you feel very unwell and can make you vomit. They can make you feel breathless. If you do not get rid of the ketones, you can become extremely unwell and would need urgent hospital treatment (diabetic ketoacidosis (DKA)).

Ketones can still be produced when you are ill even when your blood glucose is low. Therefore:

- Never stop your insulin even if you are not eating.
- Always check for ketones if you feel unwell <u>regardless of your blood</u> <u>glucose level.</u>
- Inform your Diabetes Team of illness.


To find out more about our Trust visit www.royalberkshire.nhs.uk

Please ask if you need this information in another language or format.

Paediatric Unit. Reviewed: October 2023.

Next review due: October 2025

NEVER STOP YOUR PUMP IF YOU ARE UNWELL + ALWAYS CHECK FOR KETONES IF YOU ARE UNWELL REGARDLESS OF BLOOD GLUCOSE OR IF WELL, WHEN **BLOOD GLUCOSE IS ABOVE 14 mmols/l**



CHECK BLOOD KETONES

Blood Ketones 1.0 - 2.9

Eating Not eating

Extra insulin now (see table below) + usual insulin for food given with **PEN** Change cannula, tubing and insulin

cartridge

Extra insulin now (see table below) given with PEN Change cannula, tubing and insulin cartridge

(see table below)				
+ usual insulin for				
food given with PEN				
Change cannula,				
tubing and insulin				
cartridge				

Extra insulin now

Eating

Blood Ketones 3.0 or more Inform Diabetes Team

Not eating

Extra insulin now (see table below) given with **PEN** Change cannula, tubing and insulin cartridge

ı				
l	Extra insulin			
ı	Weight	NovoRapid	Weight	NovoRapid
ı	(Kgs)	dose (units)	(Kgs)	dose (units)
l	10	2	50	10
l	20	4	60	12
l	30	6	70	14
ı	40	8	80	16

Extra insulin Weight NovoRapid Weight NovoRapid dose (units) dose (units) (Kgs) (Kgs) 10 1 50 5 20 2 60 6 70 30 3 7 40 80

Blood glucose less than 6

Frequent small sugary drinks or Glucogel

Check **Blood Glucose** Blood glucose more than 6

Frequent sugar-free drinks

Check blood ketones and glucose every 2 hours and repeat process until child is recovered

Contact team if:

Look for cause of high blood

glucose and change cannula,

tube and insulin cartridge

- Increasingly unwell
- Ketones more than 3 or persist for more than 6 hours
- Vomits more than twice
- You have to give a second extra dose of insulin
- Unable to maintain blood glucose above 4
- Worried for any reason

Phone numbers

0118 322 8922 (nurses)

After 8pm 0118 322 5111 and ask to bleep paediatric registrar