



Cervical fusion information and exercises

A cervical fusion is any operation to join two or more bones in your neck together. This can be performed, by using any combination of bone-graft, metal plates or screws.

You may be fitted with a collar, which should be worn at all times during the first 4-6 weeks.

This information is designed to help you get back to full fitness as quickly as possible after your operation.

Before you leave hospital a physiotherapist will teach you all the exercises on this leaflet and make sure that you do not have any problems.

Introduction

Everybody's symptoms are different. Those patients with arm pain only usually recover quickly and can often go home within 24 hours of surgery. In patients who have the surgery due to pressure on the spinal cord, the recovery is often slower and the symptoms don't always fully resolve.

- **Work** – return to work is dependent on the nature of your job. If you have a desk job you may feel comfortable to return at 6-8 weeks post op; a heavy manual job your return to work is likely to be after a minimum of three months. Always check with your surgeon before returning to work.
- **Driving** – never drive if you have a collar. Once the collar has been removed and you have sufficient range of movement to check the blind spots you may return to driving. If you have had cord compression resulting in weakness of the legs, you must also be able to do an emergency stop before driving.

On discharge

- Continue with your exercises as shown.
- Avoid heavy lifting for three months or as advised by your surgeon.
- If you have a collar you will be seen after the collar is removed for physiotherapy, if you have no collar you will be seen between 2 and 4 weeks after your surgery.
- You will be seen by your surgeon for review at 6-8 weeks after your operation.

Day 1 – discharge

- You will be reviewed by the physiotherapist on the ward. Your mobility will be assessed and if required you will be provided with the appropriate walking aid.
- You may mobilise freely round the ward if safe to do so.

- Start the exercises shown below, start with a minimum of twice a day. If the pain is settling and completing the exercises twice a day doesn't aggravate your symptoms, you may increase them to 3-4 times a day if you wish.
- Continue with the exercises for up to 3 months or until back to your normal activities.
- Once you are mobilising safely and have managed the stairs if you need to do so, you will be discharged home. This is can be within 24 hours of your surgery.

Exercises

1. Deep neck flexors

Within the collar, keeping head upright, pull your chin in. Hold the end position and gently nod head downwards 10 times. Relax.



2. Nerve mobilisation


These exercises are designed to keep the nerves in the arm moving and preventing them from becoming trapped in scar tissue.

3. Median nerve mobilisation

Raise your arm to shoulder height.

Start with the elbow bent then gently straighten it with the hand palm upwards. Once the elbow is straight, extend the wrist so the fingers point towards the floor.

Hold for 5 seconds, relax and then repeat 4 more times




4. Ulna nerve mobilisation

Raise your arm to shoulder height.


Bend the elbow and rotate the forearm so the palm of your hand rests against the side of your head, with your fingers pointing downwards.

Hold for 5 seconds, relax and then repeat 4 more times.



5. Radial nerve mobilisation

Keeping your arm by your side, twist the arm inwards, bend your wrist and then take the arm backwards, angling slightly away from your side. Hold for 5 seconds, relax and repeat 4 more times.




© The Saunders Group Inc.

If you have **no** soft collar you may also do the following exercises.

6. Neck flexion


Gently bend neck forward. Do not force to end of range. Relax, return to the upright position. Repeat 10 times.



© PhysioTools Ltd

7. Rotation

Gently rotate your head to one side, do not force to end of range. Rotate to the other side and relax. Repeat 10 times.



© PhysioTools Ltd

Contact us

RBFT Physiotherapy Department
0118 322 7811 or 7812

To find out more about our Trust visit www.royalberkshire.nhs.uk

Please ask if you need this information in another language or format.

RBFT Orthopaedic Physiotherapy, January 2024
Next review due: January 2026